

ON THE m -TORSION SUBGROUP OF THE BRAUER GROUP OF A GLOBAL FIELD

WEN-CHEN CHI*, HUNG-MIN LIAO, AND KI-SENG TAN†

ABSTRACT. In this note, we give a short proof of the existence of certain abelian extension over a given global field K . This result implies that for every positive integer m , there exists an abelian extension L/K of exponent m such that the m -torsion subgroup of $\text{Br}(K)$ equals $\text{Br}(L/K)$.

Assume that a global field K , a prime number p and a natural number n are given. We say that a pro- p abelian extension M/K is granted if the exponent of M/K , by which we mean that of $\text{Gal}(M/K)$, divides p^n and for every finite place w of M the degree $[M_w : K_v]$, $w|v$, also divides p^n . A finite place v is said to be supported under M/K if for each $w|v$, $[M_w : K_v] = p^n$; an archimedean place v is supported under M/K if M_w is complex for every $w|v$. The purpose of this note is to give a simple proof of the following.

Main Theorem. *Suppose K , p and n are given as above. Then there exists a granted extension that supports every place of K .*

We should remind the reader that [KS03, KS06, Po05] put together already gave a complete proof of the theorem. However, our proof is much simpler. Furthermore, as these papers have pointed out, using the structure theorem of Brauer groups over global fields, one can deduce from the main theorem the following corollary which gives an affirmative answer to the question raised in [AS02]. Recall that the relative Brauer group $\text{Br}(L/K)$ is the kernel of the restriction map $\text{Br}(K) \rightarrow \text{Br}(L)$ ([AS02]).

Corollary. *For a positive integer m , there always exists an abelian extension L/K of exponent m such that the m -torsion subgroup of $\text{Br}(K)$ equals the relative Brauer group $\text{Br}(L/K)$.*

We shall prove the theorem by constructing a sequence

$$M_0 \subset M_1 \subset \cdots \subset M_k \subset \cdots$$

of granted extensions over K and show that $L = \bigcup_{k=0}^{\infty} M_k$ enjoys the desired property. We will frequently use the fact that if an abelian extension K'/K is of exponent p^n and is unramified at v then the degree $[K'_w : K_v]$, $w|v$, divides p^n . Thus, in

* The author was supported in part by the National Science Council of Taiwan, NSC95-2115-M-003-005. † The author was supported in part by the National Science Council of Taiwan, NSC95-2115-M-002-017-MY2.

order to check if K'/K is granted, it is sufficient to check the local degrees at the ramified places.

Consider $\text{Pic}_K := K^* \backslash \mathbb{A}_K^* / \prod_v \mathcal{O}_v^*$, where $\mathbb{A}_K^*$ is the ideles group, v runs through all places of K and for an Archimedean place v we let $\mathcal{O}_v^* = K_v^*$. For a finite place $\mathfrak{p}$, let $[\mathfrak{p}]$ be the image of any prime element $\pi_{\mathfrak{p}} \in K_{\mathfrak{p}}^*$ under the natural map $K_{\mathfrak{p}}^* \rightarrow \mathbb{A}_K^* \rightarrow \text{Pic}_K$. The abelian group Pic_K is finitely generated and by the Chabotarev density theorem we can find $\mathfrak{p}_1, \mathfrak{p}_2, \dots, \mathfrak{p}_\ell$ outside any given finite set of places such that $[\mathfrak{p}_1], [\mathfrak{p}_2], \dots, [\mathfrak{p}_\ell]$ form a set of generators of Pic_K . Denote $S = \{\mathfrak{p}_1, \mathfrak{p}_2, \dots, \mathfrak{p}_\ell\}$. For later usage, we choose for every idele $x \in \mathbb{A}_K^*$ a global element $f_x \in K^*$, which is unique up to $\mathcal{O}_S^*$, such that

$$x = y \cdot f_x \cdot u \quad (0.1)$$

for some $y \in \prod_{i=1}^\ell K_{\mathfrak{p}_i}^*$ and $u \in \prod_v \mathcal{O}_v^*$.

Put $\Gamma_{-1} = \text{Pic}_K / (p^n \text{Pic}_K) = K^* \backslash \mathbb{A}_K^* / (\prod_v \mathcal{O}_v^* \cdot (\mathbb{A}_K^*)^{p^n})$. Class Field Theory ([AT90]) identifies Γ_{-1} as the Galois group of an abelian extension, denoted as M_{-1}/K , which is everywhere unramified. This together with the obvious fact that the exponent of Γ_{-1} divides p^n implies that the extension M_{-1}/K is granted.

Suppose T is a finite set of places with $T \cap S = \emptyset$ and $\mathcal{N} \subset \prod_{v \in T} \mathcal{O}_v^*$ is an open subgroup which contains $\mathcal{O}_S^*$ via the natural embedding $K^* \hookrightarrow \prod_{v \in T} K_v^*$. Set

$$\Gamma(\mathcal{N}) := K^* \backslash \mathbb{A}_K^* / ((\prod_{v \notin T} \mathcal{O}_v^* \times \mathcal{N}) \cdot (\mathbb{A}_K^*)^{p^n}).$$

This group is also of exponent dividing p^n . Again, Class Field Theory identifies $\Gamma(\mathcal{N})$ as the Galois group of an abelian extension which we denote as $M(\mathcal{N})/K$.

Lemma 0.1. *Let $\mathcal{N}$ be as above. Then we have the exact sequence*

$$0 \rightarrow \prod_{v \in T} \mathcal{O}_v^* / (\mathcal{N} \cdot \prod_{v \in T} (\mathcal{O}_v^*)^{p^n}) \xrightarrow{i} \Gamma(\mathcal{N}) \xrightarrow{q} \Gamma_{-1} \rightarrow 0, \quad (0.2)$$

where i is induced from the natural map $\prod_{v \in T} K_v^* \rightarrow \mathbb{A}_K^*$ and q is the natural quotient map.

Proof. It is enough to show the injectivity of i . Suppose $\bar{z} \in \ker(i)$ is obtained from an element $z \in \prod_{v \in T} \mathcal{O}_v^*$. Then $z = f \cdot t \cdot x^{p^n}$ for some $f \in K^*$, $t \in \prod_{v \notin T} \mathcal{O}_v^* \times \mathcal{N}$ and $x \in \mathbb{A}_K^*$. Write $x = y \cdot f_x \cdot u$ as in (0.1). Then we see that $f_x^{p^n} \cdot f \in \mathcal{O}_S^*$. Therefore if t_T and u_T are the T -components of t and u , then $z = (f_x^{p^n} \cdot f) \cdot t_T \cdot u_T^{p^n} \in \mathcal{N} \cdot (\prod_{v \in T} \mathcal{O}_v^*)^{p^n}$. Hence $\bar{z} = 0$. $\square$

Suppose $p = 2$ and $\infty_1, \dots, \infty_s$, $s > 0$, are all the real places of K . Put $\mathbb{R}_j = K_{\infty_j}$ and let sgn_j^b be the sign map $\mathbb{R}_j^* \rightarrow \mathbb{R}_j^* / \mathbb{R}_{j,+} \simeq \mathbb{Z}/2\mathbb{Z}$. As M_{-1}/K is unramified everywhere, for each j , we can choose a real place of M_{-1} sitting over ∞_j and use it to define the sign map

$$\text{sgn}_j : M_{-1}^* \longrightarrow \mathbb{R}_j^* \xrightarrow{\text{sgn}_j^b} \mathbb{Z}/2\mathbb{Z}.$$

Consider $U = \{f \in \mathcal{O}_S^* \subset K^* \mid \sum_{j=1}^s \text{sgn}_j(f) \equiv 0 \pmod{2}\}$. Then either $\mathcal{O}_S^* = U$ or $\mathcal{O}_S^* = U \amalg gU$ for some g . Let M'_{-1} be the composite of all $M_{-1}(\sqrt{f})$, $f \in U$. By the Chabotarev density theorem we can find a finite place $\mathfrak{p}$ splitting completely

under M'_{-1}/K . Since on $(M_{-1}^*)^2$ all the values of the map $\text{sgn} := \sum_{j=1}^s \text{sgn}_j$ equal $0 \in \mathbb{Z}/2\mathbb{Z}$, if g exists then $U \cdot (M_{-1}^*)^2 \subsetneq \mathcal{O}_S^* \cdot (M_{-1}^*)^2$. In this case, Kummer's Theory says that $M'_{-1}(\sqrt{g})/M'_{-1}$ is a quadratic extension and we choose $\mathfrak{p}$ so that it does not completely split under $M'_{-1}(\sqrt{g})/K$.

Let $\text{sgn}_{\mathfrak{p}}^b$ be the map $\mathcal{O}_{\mathfrak{p}}^* \rightarrow \mathcal{O}_{\mathfrak{p}}^*/(\mathcal{O}_{\mathfrak{p}}^*)^2 \simeq \mathbb{Z}/2\mathbb{Z}$, for $(x_{\mathfrak{p}}, x_1, \dots, x_s) \in \mathcal{O}_{\mathfrak{p}}^* \times \prod_{i=1}^s \mathbb{R}_i^*$ define $\text{sgn}^b(x_{\mathfrak{p}}, x_1, \dots, x_s) = \text{sgn}_{\mathfrak{p}}^b(x_{\mathfrak{p}}) + \sum_{i=1}^s \text{sgn}_i^b(x_i)$ and put

$$\mathcal{N}_0 = \{(x_{\mathfrak{p}}, x_1, \dots, x_s) \in \mathcal{O}_{\mathfrak{p}}^* \times \prod_{i=1}^s \mathbb{R}_i^* \mid \text{sgn}^b(x_{\mathfrak{p}}, x_1, \dots, x_s) \equiv 0 \pmod{2}\}.$$

The extension $M(\mathcal{N}_0)/K$ is unramified outside $\{\mathfrak{p}, \infty_1, \dots, \infty_s\}$. Since each $\mathbb{R}_i^* \not\subset \mathcal{N}_0$, the extension is ramified at each ∞_i . The map sgn^b induces an isomorphism $\mathcal{O}_{\mathfrak{p}}^* \times \prod_{i=1}^s \mathbb{R}_i^*/\mathcal{N}_0 \simeq \mathbb{Z}/2\mathbb{Z}$ and from Lemma 0.1 we see that (since $\mathfrak{p}$ splits completely under M'_{-1}/K) the local extension of $M(\mathcal{N}_0)/K$ at $\mathfrak{p}$ is a quadratic extension. Therefore $M(\mathcal{N}_0)/K$ is granted and it supports every real place. We put $M_0 = M(\mathcal{N}_0)$ if $p = 2$ and K has a real place. Otherwise, put $M_0 = M_{-1}$.

We set all finite places of K into a sequence $\mathfrak{q}_1, \mathfrak{q}_2, \dots, \mathfrak{q}_k, \dots$. And we shall construct M_k so that it supports $\mathfrak{q}_k$. Assume that M_{k-1} is already constructed. If it supports $\mathfrak{q}_k$, then we set $M_k = M_{k-1}$. Otherwise, assume that the decomposition subgroup of $\text{Gal}(M_{k-1}/K)$ at $\mathfrak{q}_k$ is of order p^m with $m < n$. We have $m \geq m_1 + m_2$ where p^{m_1} is the order of the inertia subgroup of $\text{Gal}(M_{k-1}/K)$ and p^{m_2} is the degree of the residue field extension of M_{-1}/K at $\mathfrak{q}_k$.

Let $S_1 = \{v_1, \dots, v_r\}$ be a finite set of places of K such that $\mathfrak{q}_k \notin S_1$ and M_{k-1}/K is unramified outside $S_1 \cup \{\mathfrak{q}_k\}$. We can assume that S is chosen so that $S \cap (S_1 \cup \{\mathfrak{q}_k\}) = \emptyset$. And denote $S_2 = S \cup S_1$, $S_3 = S_2 \cup \{\mathfrak{q}_k\}$. Choose a prime element π at $\mathfrak{q}_k$ and let $f_{\pi} \in K^*$ be the global element chosen before. Then the S_3 -units group $\mathcal{O}_{S_3}^*$ is the direct product of $\mathcal{O}_{S_2}^*$ and the infinite cyclic subgroup generated by f_{π} .

To construct M_k , we will need to find a finite place w outside S_3 , which splits completely under M_{k-1}/K , and an open subgroup $\mathcal{N}_w \subset \mathcal{O}_w^*$, which contains $\mathcal{O}_{S_2}^*$ via the natural embedding $K^* \hookrightarrow K_w^*$, such that the quotient $\mathcal{O}_w^*/\mathcal{N}_w$ is isomorphic to $\mathbb{Z}/p^n\mathbb{Z}$ and contains a subgroup of order p^{n-m_1} generated by the element $f_{\pi} \pmod{\mathcal{N}_w}$.

Lemma 0.2. *If $\mathcal{N}_w$ is as above, then the following hold:*

- (1) *The extension $M(\mathcal{N}_w)/K$ is unramified outside $\{w\}$.*
- (2) *Every finite place $\mathfrak{q} \in S_2$ splits completely under the extension $M(\mathcal{N}_w)/M_{-1}$.*
- (3) *If $\mathfrak{q} \mid \mathfrak{q}_k$, then the degree $[M(\mathcal{N}_w)_{\mathfrak{q}} : K_{\mathfrak{q}_k}] = p^{n-m_1}$.*
- (4) *The extension is totally ramified at w with $[M(\mathcal{N}_w)_w : K_w] = p^n$.*

Proof. Statements (1) and (4) are from the exact sequence (0.2), since the place w splits completely under M_{-1}/K . To prove (2), let $\pi_{\mathfrak{q}}$ be a prime element at $\mathfrak{q}$ and assume that in Γ_{-1} the Frobenius at $\mathfrak{q}$ is annihilated by p^{μ} . Then $\pi_{\mathfrak{q}}^{p^{\mu}} = f \cdot t \cdot x^{p^n}$ for some $f \in K^*$, $t \in \prod_v \mathcal{O}_v^*$ and $x \in \mathbb{A}_K^*$. This and the equation (0.1) imply that $f \cdot f_x^{p^n} \in \mathcal{O}_{S_2}^*$. They also imply that t_w , the w -component of t , is contained in $\mathcal{N}_w$, since $\mathcal{O}_{S_2}^* \cdot (\mathcal{O}_w^*)^{p^n} \subset \mathcal{N}_w$. Therefore in $\Gamma(\mathcal{N}_w)$ the Frobenius at $\mathfrak{q}$ is also annihilated by p^{μ} . And (2) is proved.

Let $F_r \in \Gamma(\mathcal{N}_w)$ be the Frobenius element at $\mathfrak{q}_k$. Using the exact sequence (0.2), we deduce that $p^{m_2}F_r \in \mathcal{O}_w^*/\mathcal{N}_w$ and hence $\pi^{m_2} = f \cdot t \cdot x^{p^n}$ for some $f \in K^*$, $t \in \prod_v \mathcal{O}_v^*$ and $x \in \mathcal{A}_K^*$. This means $p^{m_2}F_r$ equals the residue class $t_w \pmod{\mathcal{N}_w}$ and, together with the equation (0.1), this also shows that $t_w \equiv f^{-1} \pmod{(\mathcal{O}_w^*)^{p^n}}$. We then write $\pi = z \cdot f_\pi \cdot u'$ with $z \in \prod_{i=1}^s K_{p_i}^*$ and $u' \in \prod_v \mathcal{O}_v^*$ and use it to deduce that $f^{-1} \cdot f_x^{-p^n} \cdot f_\pi^{p^{m_2}} \in \mathcal{O}_{S_2}^*$. Again, since $\mathcal{O}_{S_2}^* \cdot (\mathcal{O}_w^*)^{p^n} \subset \mathcal{N}_w$, we have $t_w \equiv f_\pi^{p^{m_2}} \pmod{\mathcal{N}_w}$. Therefore in the quotient $\mathcal{O}_w/\mathcal{N}_w$ the order of the residue class $t_w \pmod{\mathcal{N}_w}$ equals $p^{n-m_1-m_2}$. And this proves that F_r is of order p^{n-m_1} . $\square$

We then set $M_k = M_{k-1}M(\mathcal{N}_w)$. It is clear that the exponent of $\text{Gal}(M_k/K)$ is p^n . The extension M_k/K is unramified outside $S_1 \cup \{q_k, w\}$. Lemma 0.2 implies that $[M_{k,\mathfrak{p}} : K_{\mathfrak{q}_k}] = p^n$ for $\mathfrak{p} \mid \mathfrak{q}_k$, $[M_w : K_w] = p^n$ and if $\mathfrak{q} \in S_1$, then $[M_{k,\mathfrak{q}} : K_{\mathfrak{q}}] = [M_{k-1,\mathfrak{q}} : K_{\mathfrak{q}}]$. Therefore M_k/K is granted and it supports $\mathfrak{q}_k$.

To complete the proof, we need to find w and $\mathcal{N}_w$. Let us first consider the case where $\text{char.}(K) \neq p$. Let $\mathfrak{q}$ be a place of M_{k-1} sitting over $\mathfrak{q}_k$. Then we have $\text{ord}_{\mathfrak{q}}(f_\pi) = p^{m_1} \cdot \text{ord}_{\mathfrak{q}_k}(f_\pi) = p^{m_1}$. Since $\text{ord}_{\mathfrak{q}}(f) = 0$ for all $f \in \mathcal{O}_{S_2}^*$, the index $|\mathcal{O}_{S_2}^* \cdot (M_{k-1}^*)^{p^n} : \mathcal{O}_{S_3}^* \cdot (M_{k-1}^*)^{p^n}|$ is a multiple of p^{n-m_1} . For a global element $f \in M_{k-1}^*$ let $M_{k-1}(\sqrt[p^n]{f})$ be the Kummer extension generated by all p^n th root of f . Let K'_2 be the composite of all $M_{k-1}(\sqrt[p^n]{f})$, $f \in \mathcal{O}_{S_2}^*$ and let K'_3 be the composite of all $M_{k-1}(\sqrt[p^n]{f})$, $f \in \mathcal{O}_{S_3}^*$. Kummer's theory tells us that the degree $[K'_3 : K'_2]$ is a multiple of p^{n-m_1} . Choose an element $\sigma \in \text{Gal}(K'_3/K'_2) \subset \text{Gal}(K'_3/K)$ of order p^{n-m-1} and apply the Chabotarev density theorem to choose w to be a unramified place under K'_3/K such that the Frobenius at w equals σ . We choose w outside S_3 and put $\mathcal{N}_w = (\mathcal{O}_w^*)^{p^n}$. It is obvious that w splits completely under M_{k-1} , $\mathcal{O}_{S_2}^* \subset \mathcal{N}_w$ and in the quotient $\mathcal{O}_w^*/\mathcal{N}_w$ the order of $f_\pi \pmod{\mathcal{N}_w}$ equals p^{n-m_1} . Since K'_2 contains all the primitive roots of 1 and w splits completely under K'_2/K , the local ring $\mathcal{O}_w$ also contains all the primitive roots of 1. This implies that $\mathcal{O}_w^*/\mathcal{N}_w \simeq \mathbb{Z}/p^n\mathbb{Z}$. Thus w and $\mathcal{N}_w$ satisfy all the required conditions.

Finally, we consider the case where $\text{char.}(K) = p$. Apply the Chabotarev density theorem and choose w to be a place outside S_2 , splitting completely under M_{k-1}/K . Recall that the p -part $\mathcal{O}_1 = 1 + \pi_w \mathcal{O}_w$ of $\mathcal{O}_w^*$ is the direct product of countable many copies of $\mathbb{Z}_p$ ([We67]) and the local Leopoldt Conjecture holds ([Kis93]) in the sense that $\mathbb{Z}_p \otimes_{\mathbb{Z}} \mathcal{O}_{S_3}^*$ form a direct summand of $\mathcal{O}_1$. In other words, we have

$$\mathcal{O}_1 = \mathbb{Z}_p \otimes \mathcal{O}_{S_2}^* \times \mathbb{Z}_p \otimes C \times W$$

where C is the infinite cyclic group generated by f_π and W is a direct product of countable many copies of $\mathbb{Z}_p$. Using these, we can easily find an open subgroup $\mathcal{N}_w \subset \mathcal{O}_w^*$ so that $\mathcal{O}_w^*/\mathcal{N}_w \simeq \mathbb{Z}/p^n\mathbb{Z}$, $f \in \mathcal{N}_w \forall f \in \mathcal{O}_{S'}^*$ and f_π generates a subgroup of order p^{n-m_1} in $\mathcal{O}_w^*/\mathcal{N}_w$.

REFERENCES

- [AS02] E. Aljadeff and J. Sonn, *Relative Brauer groups and m -torsion*, Proc. Amer. Math. Soc. 130(2002), 1333-1337.
- [AT90] E. Artin and J. Tate, *Class Field Theory*, Advanced Book Classics, Vol.47, Addison-Wesley Publishing Co., Inc., Reading, MA, 1990.
- [Iw86] K. Iwasawa, *Local Class Field Theory*, Oxford University Press, 1986.

- [Kis93] Kisilevsky, H.: *Multiplicative independence in function fields*. J. Number Theory **44**, 352–355 (1993)
- [KS03] H. Kisilevsky and J. Sonn, *On the n -torsion subgroup of the Brauer group of a number field*, J. Theor. Nombres Bordeaux 15(2003), 199-204.
- [KS06] H. Kisilevsky and J. Sonn, *Abelian extensions of global fields with constant local degrees*, Math. Res. Letters.
- [Po05] C. D. Popescu, *Torsion subgroups of Brauer groups and extensions of constant local degree for global function fields*, J. Number Theory 115(2005), 27-44.
- [We67] A. Weil, *Basic Number theory*, Springer-Verlag, Berlin-New York, 1967.

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN NORMAL UNIVERSITY, 88, SEC.4, TING-CHOU ROAD, TAIPEI, TAIWAN 116, E-MAIL: WCHI@MATH.NTNU.EDU.TW

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN NORMAL UNIVERSITY, 88, SEC.4, TING-CHOU ROAD, TAIPEI, TAIWAN 116, E-MAIL: 89340001@NTNU.EDU.TW

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, 1, SEC.4, ROOSEVELT ROAD, TAIPEI, TAIWAN 106, E-MAIL: TAN@MATH.NTU.EDU.TW